

Many-to-many disjoint paths in hypercubes with faulty vertices ^{*}

Xiang-Jun Li^a Bin Liu^a Meijie Ma^b Jun-Ming Xu^{a†}

^aSchool of Mathematical Sciences

University of Science and Technology of China

Wentsun Wu Key Laboratory of CAS

Hefei 230026, China

^bDepartment of Mathematics

Zhejiang Normal University

Jinhua 321004, China

Abstract

This paper considers the problem of many-to-many disjoint paths in the hypercube Q_n with f faulty vertices and obtains the following result. For any integer k with $1 \leq k \leq n - 2$, any two sets S and T of k fault-free vertices in different parts of Q_n ($n \geq 3$), if $f \leq 2n - 2k - 3$ and each fault-free vertex has at least two fault-free neighbors, then there exist k fully disjoint fault-free paths linking S and T which contain at least $2^n - 2f$ vertices. This result improves some known results in a sense.

Keywords: hypercube, disjoint paths, fault-tolerance, parallel computing

AMS Subject Classification (2000): 05C38, 05C45, 68M10, 68M15, 68R10, 90B10

1 Induction

The n -dimensional hypercube Q_n is a graph whose vertex-set consists of all binary vectors of length n , with two vertices being adjacent whenever the corresponding vectors differ in exactly one coordinate. There is a large amount of literature on graph-theoretic properties of hypercubes (e.g., see the comprehensive survey paper on early results [7])

^{*}The work was supported by NNSF of China (Nos 10711233, 11101378) and ZSDZZZZXK08.

[†]Corresponding author: xujm@ustc.edu.cn (J.-M. Xu)

and recent results [13] as well as on their applications in parallel computing (e.g., see [10]).

One of most central issues in various high performance communication networks or parallel computing systems is to find a cycle or a path of given length in Q_n (see [5, 6, 8] and the survey paper [15]). To find node-disjoint paths concerned with fault-tolerant routings among nodes has received much research attention recently (see, for example, [3, 11, 12] and references cited therein). In this paper, we consider the problem of node disjoint paths of Q_n .

It is well known that there are n internally disjoint paths between any two vertices u and v in Q_n , but they do not always contain all vertices. In 2004, Chang *et al.* [1] showed that for any integer k with $1 \leq k \leq n$ and any two vertices u, v from different partite sets of Q_n , there are k internally disjoint uv -paths in Q_n which contains all vertices. In 2007, Chang *et al.* [2] further showed that for any integer k with $1 \leq k \leq n - 4$ and any two vertices u, v from different partite sets of Q_n ($n \geq 5$), there are k internally disjoint uv -paths in Q_n which contain all vertices and the difference of lengths of any two paths is most two. In 2009, Chen [3] showed that for any integer k with $1 \leq k \leq n - 1$, any two sets S and T of k fault-free vertices in different partite sets in Q_n with f faulty vertices and h faulty edges, if $f + h \leq n - k - 1$, then there exist k disjoint fault-free paths linking S and T in Q_n which contain at least $2^n - 2f$ vertices. In this paper, we obtain the following results.

Theorem *Let Q_n ($n \geq 3$) be an n -dimensional hypercube with f faulty vertices, and let k be an integer with $1 \leq k \leq n - 2$. If $f \leq 2n - 2k - 3$ and each fault-free vertex has at least two fault-free neighbors then, for any two sets S and T of k fault-free vertices in different partite sets, there exist k disjoint fault-free paths linking S and T in Q_n which contain at least $2^n - 2f$ vertices.*

Clearly, our theorem partly improves the above-mentioned results. The proof of our result is in Section 3. Section 2 gives some notations and lemmas.

2 Notations and Lemmas

For graph-theoretical terminology and notation not defined here, we follow [14]. Let $G = (V, E)$ be a connected simple graph, where $V = V(G)$ is the vertex-set and $E = E(G)$ is the edge-set of G . For $uv \in E(G)$, we call u (resp. v) is a neighbor of v (resp. u). A uv -path is a sequence of adjacent vertices, written as $\langle v_0, v_1, v_2, \dots, v_m \rangle$, in which $u = v_0, v = v_m$ and all the vertices $v_0, v_1, v_2, \dots, v_m$ are different from each other. The length of a path P is the number of edges in P . Let $d_G(u, v)$ be the length of a shortest uv -path in G , called the distance between u and v in G . For a path $P = \langle v_0, v_1, \dots, v_i, v_{i+1}, \dots, v_m \rangle$, we can write $P = P(v_0, v_i) + v_i v_{i+1} + P(v_{i+1}, v_m)$, and the notation $P - v_i v_{i+1}$ denotes the subgraph obtained from P by deleting the edge $v_i v_{i+1}$. Two paths are disjoint if they have no vertices in common. Given two disjoint sets S and T of k vertices, if there exist k disjoint paths between S and T , we call these paths to be k disjoint ST -paths.

The n -dimensional hypercube Q_n is a graph with 2^n vertices, each vertex denoted by an n -bit binary string $u = u_n u_{n-1} \cdots u_2 u_1$. Two vertices are adjacent if and only if their strings differ in exactly one bit position. It has been proved that Q_n is a vertex- and edge-transitive bipartite graph.

By definition, for any $k \in \{1, 2, \dots, n\}$, Q_n can be expressed as $Q_n = L_k \odot R_k$, where L_k and R_k are two subgraphs of Q_n induced by the vertices with the k bit position is 0 and 1, respectively, which are isomorphic to Q_{n-1} , linking by an edge a vertex in L_k to a vertex in R_k if they differ in only the k -th bit. Without loss of generality, we write $Q_n = L \odot R$. For convenience, for a vertex u in L , we use u_R to denote its only neighbor in R . Similarly, for a vertex v in R , we use v_L to denote its only neighbor in L . Clearly, for any two vertices u_L and v_L in L , $d_L(u_L, v_L) = d_R(u_R, v_R)$.

Let F denote a set of faulty vertices in Q_n , and $f = |F|$. When $Q_n = L \odot R$, we denote $f_L = |F \cap L|$ and $f_R = |F \cap R|$. A subgraph of Q_n is *fault-free* if it contains no vertices in F . For two subsets A, B of $V(Q_n)$, let $d(A, B) = \min\{d(x, y) : x \in A, y \in B\}$.

Lemma 2.1 (Kueng *et al.* [9], 2009) *If $f \leq 2n - 5$ and each fault-free vertex of Q_n ($n \geq 3$) has at least two fault-free neighbors then, for any two distinct fault-free vertices x and y with distance d , there is a fault-free xy -path containing at least $2^n - 2f$ vertices if d is odd and $2^n - 2f - 1$ vertices if d is even.*

Lemma 2.2 (Chen [3], 2009) *For any integer k with $1 \leq k \leq n - 1$, if $f \leq n - k - 1$ then, for any two sets S and T of k fault-free vertices in different partite sets in Q_n ($n \geq 2$), there exist k disjoint fault-free ST -paths which contain at least $2^n - 2f$ vertices.*

Lemma 2.3 (Dvořák [4], 2005) *Let (x, y) and (u, v) be two disjoint pairs of vertices with odd distance in Q_n . If x and y are adjacent in Q_n with $n \geq 3$, then there exists a uv -path containing all vertices in $Q_n - \{x, y\}$ unless $n = 3, d(u, v) = 1$ and $d(\{x, y\}, \{u, v\}) = 2$.*

Lemma 2.4 *Let x and y be two adjacent vertices of Q_n ($n \geq 4$). Then, for any integer k with $1 \leq k \leq n - 2$ and for any two sets S and T of k vertices in different partite sets in $Q_n - \{x, y\}$, there exist k disjoint ST -paths containing all vertices in $Q_n - \{x, y\}$.*

Proof. By induction on k . If $k = 1$, the result is true by Lemma 2.3. We assume that the result holds for any integer fewer than k , and consider the case of $k (\geq 2)$. Let $\{X, Y\}$ be a bipartition of Q_n and $Q_n = L \odot R$ such that the edge $xy \in E(L)$, $S = \{s_1, s_2, \dots, s_k\}$ and $T = \{t_1, t_2, \dots, t_k\}$ be two sets of k vertices in different partite sets in $Q_n - \{x, y\}$, and let

$$\begin{aligned} S_L &= S \cap L, \quad T_L = T \cap L \quad \text{and} \quad p = |S_L|, \quad q = |T_L|, \\ S_R &= S \cap R, \quad T_R = T \cap R. \end{aligned}$$

Without loss of generality, we can assume $S \cup \{x\} \subseteq X$, $T \cup \{y\} \subseteq Y$, $p \geq q$ and $0 \leq q \leq k-1$. If $p > q$, let

$$\begin{aligned} U_L &= \{u_1, \dots, u_{p-q}\} \subseteq Y \cap V(L - (T_L \cup \{y\})), \\ U_R &= \{v_1, \dots, v_{p-q}\} \subseteq X \cap V(R) \text{ with } u_i v_i \in E(Q_n) \text{ for } 1 \leq i \leq p-q. \end{aligned}$$

If $p = q$, let $U_L = U_R = \emptyset$.

The process of our induction steps strongly depends on Lemma 2.3. When $n = 4$, $L \cong R \cong Q_3$, while in Q_3 , we have to consider some exceptional cases when we apply Lemma 2.3. Thus, we first prove the case of $n \geq 5$ by considering the following three cases.

Case 1. $p = 0$. In this case, both S and T are in R . By Lemma 2.2, there exist k disjoint ST -paths $P_1, P_2, \dots, P_k$ containing all vertices in R . We can choose an edge $u_L v_L$ in $L - \{x, y\}$ such that $u_R v_R$ is an edge in some path P_i . Since $n \geq 5$, by Lemma 2.3, there exists a $u_L v_L$ -path P_0 containing all vertices in $L - \{x, y\}$. Let

$$P'_i = P_i - u_R v_R + u_R u_L + v_L v_R + P_0.$$

Replacing P_i by P'_i obtains k disjoint ST -paths $P_1, \dots, P_{i-1}, P'_i, P_{i+1}, \dots, P_k$ as required.

Case 2. $1 \leq p \leq k-1$. Let $T'_L = T_L \cup U_L$ and $S'_R = S_R \cup U_R$. Then $|T'_L| = q + (p - q) = p = |S_L|$ and $|S'_R| = (k - p) + (p - q) = k - q = |T_R|$. Since $n \geq 5$, by the induction hypothesis, there are p disjoint $S_L T'_L$ -paths containing all vertices in $L - \{x, y\}$, in which let $P_1, P_2, \dots, P_q$ be $S_L T_L$ -paths, and $P'_1, P'_2, \dots, P'_{p-q}$ be $S_L U_L$ -paths, without loss of generality, say P'_i connecting one vertex in S_L to u_i . By Lemma 2.2, there are $k - q$ disjoint $S'_R T_R$ -paths containing all vertices in R , in which let $P''_1, P''_2, \dots, P''_{p-q}$ be $U_R T_R$ -paths, say P''_i connecting v_i to one vertex in T_R , and $P_{p+1}, P_{p+2}, \dots, P_k$ be $S_R T_R$ -paths. Let

$$P_{q+i} = P'_i + u_i v_i + P''_i \text{ for each } i = 1, 2, \dots, p - q.$$

Then $P_1, P_2, \dots, P_k$ are k disjoint ST -paths containing all vertices in $Q_n - \{x, y\}$.

Case 3. $p = k$. By the hypothesis of $q \leq k-1$, $T_R \neq \emptyset$, say $t_k \in T_R$. Let $s_k \in S$, $S' = S - \{s_k\}$, $U' = U_L - \{u_{k-q}\}$ and $T' = T_L \cup U'$. Since $n \geq 5$, by the induction hypothesis, there exist $k-1$ disjoint $S' T'$ -paths containing all vertices in $L - \{x, y\}$, in which, let $P_1, P_2, \dots, P_q$ be $S' T_L$ -paths, and $P'_1, P'_2, \dots, P'_{k-1-q}$ be $S' U'$ -paths, where P'_i connects one vertex of S' to u_i for each $i = 1, 2, \dots, k-1-q$.

Assume that s_k is in some path connecting s_i to some vertex t' in T' . Let u_L be the neighbor of s_k in the path closer s_i and u_R be the neighbor of u_L in R , and let $u_R = v_{k-q}$. By Lemma 2.2, there are $k-q$ disjoint $U_R T_R$ -paths $P''_1, P''_2, \dots, P''_{k-1-q}, P''_{k-q}$ that contain all vertices in R . Without loss of generality, assume that P''_i connects v_i and t_{q+i} for each $i = 1, 2, \dots, k-q$. Let

$$P_{q+i} = P'_i + u_i v_i + P''_i \text{ for each } i = 1, 2, \dots, k-1-q.$$

If $t' \in T_L$, say $t' = t_1$ and s_k in P_1 connecting s_1 to t_1 (see Fig. 1 (a)), let

$$\begin{aligned} P_1^* &= P_1(s_k, t_1), \\ P_k &= P_1(s_1, u_L) + u_L u_R + P''_{k-q}. \end{aligned}$$




Figure 1: Illustrations for the proof of Case 3 of Lemma 2.4.

Then $P_1^*, P_2, \dots, P_{k-1}, P_k$ are k disjoint ST -paths containing all vertices in $Q_n - \{x, y\}$.

If $t' \in U'$, say $t' = u_1$ and s_k in P'_{q+1} connecting s_{q+1} to u_1 (see Fig. 1 (b)), let

$$\begin{aligned} P_{q+1}^* &= P'_{q+1}(s_k, u_1) + u_1 v_1 + P''_{q+1}, \\ P_k &= P'_{q+1}(s_1, u_L) + u_L u_R + P''_{k-q}. \end{aligned}$$

Then $P_1, \dots, P_q, P_{q+1}^*, P_{q+2}, \dots, P_{k-1}, P_k$ are k disjoint ST -paths containing all vertices in $Q_n - \{x, y\}$.

Summing up the three cases, we prove the lemma when $n \geq 5$.

When $n = 4$, we have $k = 2$. In this case, if there exists i ($1 \leq i \leq 2$) such that $d(\{s_i, t_i\}, \{x, y\}) = 2$, we can choose $Q_4 = L \odot R$ such that $xy \in E(L)$ and $\{s_i, t_i\} \subseteq R$. Then, by Lemma 2.3, we can direct verify that the lemma holds, and the details are omitted. $\blacksquare$

3 Proof of Theorem

The proof proceeds by induction on $n \geq 3$. If $k = 1$, then $f \leq 2n - 5$. The theorem follows by Lemma 2.1. If $k = n - 2$, then $f \leq 1$. The theorem follows by Lemma 2.2. Thus, the theorem holds for $3 \leq n \leq 4$. In the following discussion, we assume $n \geq 5$ and $2 \leq k \leq n - 3$.

Let Q_n be an n -dimensional hypercube, $\{X, Y\}$ be a bipartition of $V(Q_n)$, $S = \{s_1, s_2, \dots, s_k\} \subset X$ and $T = \{t_1, t_2, \dots, t_k\} \subset Y$ be any two sets of k fault-free vertices. Our aim is to construct k disjoint fault-free ST -paths containing at least $2^n - 2f$ vertices.

Since $k \geq 2$, we have $f \leq 2n - 7$. It is easy to see that there is some $j \in \{1, 2, \dots, n\}$ such that each fault-free vertex in L_j and R_j has at least two fault-free neighbors. Let

$Q_n = L \odot R$, where $L = L_j$ and $R = R_j$. Let

$$\begin{aligned} S_L &= S \cap L, \quad T_L = T \cap L \quad \text{and} \quad p = |S_L|, \quad q = |T_L|, \\ S_R &= S \cap R, \quad T_R = T \cap R. \end{aligned}$$

By symmetry, we may assume $p \geq q$.

We construct k required ST -paths by considering three cases.

Case 1. $q = k$ or $p = 0$.

In this case, two sets S and T both are in L or R . By symmetry, we consider the former one, that is, S and T both are in L . There are two subcases.

Subcase 1.1. $f_L \leq 2n - 2k - 5$.

Since $f_L \leq 2n - 2k - 5 = 2(n - 1) - 2k - 3$ and $k \leq n - 3 = (n - 1) - 2$, by the induction hypothesis, in L there are k disjoint ST -paths

$$P'_1, P'_2, \dots, P'_k \tag{3.1}$$

containing at least $2^{n-1} - 2f_L$ vertices. Note that, when $n \geq 5$ and $f \leq 2n - 7$,

$$\begin{aligned} \sum_{i=1}^k |E(P'_i)| &= \sum_{i=1}^k |V(P'_i)| - k \\ &\geq 2^{n-1} - 2f_L - k \\ &\geq 2^{n-1} - 2f - (n - 3) \\ &> 4(2n - 7) - 2f \\ &\geq 2f. \end{aligned}$$

Therefore, there is an edge $u_L v_L$ in some path, say P'_1 , such that u_R and v_R both are fault-free. Since $f_R \leq f \leq 2(n - 1) - 5$, by Lemma 2.1, in R there is an $u_R v_R$ -path P_R containing at least $2^{n-1} - 2f_R$ vertices. Let

$$P_1 = P'_1 - u_L v_L + u_L u_R + P_R + v_R v_L, \quad P_i = P'_i \quad \text{for each } i = 2, 3, \dots, k. \tag{3.2}$$

Then $P_1, P_2, \dots, P_k$ are k disjoint ST -paths containing at least $2^n - 2f$ vertices in Q_n .

Subcase 1.2. $f_L \geq 2n - 2k - 4$.

In this case, $f_R \leq 1$ since $f \leq 2n - 2k - 3$. We consider two subcases according as whether f_R is equal to 1 or not, respectively.

Subcase 1.2a. $f_R = 1$.

In this case, $f = 2n - 2k - 3$ and $f_L = 2n - 2k - 4$. All paths in (3.1) contain at most one faulty vertex. If they contain no faulty vertex, then paths defined in (3.2) are as required. Assume that some path, say P'_1 , contains a faulty vertex w . Without loss of generality, P'_1 connects s_1 and t_1 . Note that, in this case, paths in (3.1) contain at least $2^{n-1} - 2(f_L - 1)$ vertices in L . Let u_L and v_L be two neighbors of w in P'_1 , where u_L is in $P'_1(s_1, w)$ and v_L is in $P'_1(w, t_1)$. Then the distance of u_L and v_L is even.

If u_R and v_R both are fault-free then, by Lemma 2.1, there is a fault-free $u_R v_R$ -path P_R containing at least $2^{n-1} - 2f_R - 1$ vertices in R . Let

$$P_1^* = P'_1(s_1, u_L) + u_L u_R + P_R + v_R v_L + P'_1(v_L, t_1).$$

Then, replacing P_1 in (3.2) by P_1^* yields k disjoint fault-free ST -paths with at least $2^n - 2f$ vertices in Q_n .

If one of u_R and v_R is faulty vertex, say u_R , then u_R is the only faulty vertex in R . If $u_L \neq s_1$, let z_L be the fault-free neighbor of u_L in $P'_1(s_1, u_L)$ (maybe $z_L = s_1$), then z_R is fault-free in R , and the distance z_R and v_R is odd. By Lemma 2.1, there is a fault-free $z_R v_R$ -path P_R containing at least $2^{n-1} - 2f_R$ vertices in R . Let

$$P_1^* = P'_1(s_1, z_L) + z_L z_R + P_R + v_R v_L + P'_1(v_L, t_1).$$

Replacing P_1 in (3.2) by P_1^* yields k disjoint fault-free ST -paths with vertices at least

$$(2^{n-1} - 2(f_L - 1) - |\{w, u_L\}|) + (2^{n-1} - 2f_R) = 2^n - 2f.$$

We now assume $u_L = s_1$ and u_R is the only faulty vertex in R . Note that u_L has at least two fault-free neighbor in L .

If there is a fault-free neighbor z_L of u_L not in P'_i for each $i = 1, 2, \dots, k$, we consider two vertices z_R and v_R in R . They are free-fault in R since u_R is the only faulty vertex in R . Since $d(z_L, v_L) = 3$, the distance between z_R and v_R is odd. Let P_R be a $z_R v_R$ -path with at least $2^{n-1} - 2f_R$ vertices in R , and let

$$P_1^* = s_1 z_L + z_L z_R + P_R + v_R v_L + P'_1(v_L, t_1).$$

Replacing P_1 in (3.2) by P_1^* yields k disjoint fault-free ST -paths with vertices at least

$$(2^{n-1} - 2(f_L - 1) - |\{w\}|) + (2^{n-1} - 2f_R) > 2^n - 2f.$$

If all the fault-free neighbors of u_L are in P'_1 , then one of them, say z_L , is not t_1 . Clearly, $z_L \neq v_L$ since Q_n contains no triangles. Let z'_L be the neighbor of z_L in $P'_1(z_L, t_1)$ (maybe $z'_L = t_1$), then z'_R is fault-free in R . Since both v_L and z'_L are in X , the distance between v_R and z'_R is even. Let P_R be a $v_R z'_R$ -path with at least $2^{n-1} - 2f_R - 1$ vertices in R , and let

$$P_1^* = s_1 z_L + P'_1(z_L, v_L) + v_L v_R + P_R + z'_R z'_L + P'_1(z'_L, t_1).$$

Replacing P_1 in (3.2) by P_1^* yields k disjoint fault-free ST -paths with vertices at least

$$(2^{n-1} - 2(f_L - 1) - |\{w\}|) + (2^{n-1} - 2f_R - 1) = 2^n - 2f.$$

If there is a fault-free neighbor z_L of u_L not in P'_1 , without loss of generality, assume that z_L is in P'_2 , and P'_2 connect s_2 and t_2 (see Fig. 2). Then $z_L \neq s_2$ since $z_L \in Y$ and $s_2 \in X$. Let z'_L be the neighbor of z_L in $P'_2(s_2, z_L)$ (maybe $z'_L = s_2$), then z'_R is fault-free in R . Since both v_L and z'_L are in X , both v_R and z'_R are in Y , and so the




Figure 2: An illustrations for the proof of Subcase 1.2a of the theorem.

distance between z'_R and v_R is even. Let P_R be a $z'_R v_R$ -path with at least $2^{n-1} - 2f_R - 1$ vertices in R , and let

$$P_1^* = s_1 z_L + P_2'(z_L, t_2), \quad P_2^* = P_2'(s_2, z'_L) + z'_L z'_R + P_R + v_R v_L + P_1'(v_L, t_1).$$

Replacing P_1 and P_2 in (3.2) by P_1^* and P_2^* yields k disjoint fault-free ST -paths with vertices at least

$$(2^{n-1} - 2(f_L - 1) - |\{w\}|) + (2^{n-1} - 2f_R - 1) = 2^n - 2f.$$

Summing up the above discussion, the theorem holds if $f_R = 1$.

Subcase 1.2b. $f_R = 0$.

Let $S' = S - \{s_k\}$ and $T' = T - \{t_k\}$. Since $f_L = f \leq 2n - 2k - 3 = 2(n-1) - 2(k-1) - 3$, by the induction hypothesis, in L there are $(k-1)$ disjoint fault-free $S'T'$ -paths

$$P'_1, P'_2, \dots, P'_{k-1} \quad (3.3)$$

containing at least $2^{n-1} - 2f$ vertices. Let

$$K = V(P'_1 \cup P'_2 \cup \dots \cup P'_{k-1}),$$

and let s_R and t_R be neighbors of s_k and t_k in R , respectively. Since s_k and t_k are in different partite sets in Q_n , $d_L(s_k, t_k)$ is odd, and so is $d_R(s_R, t_R)$.

If $|\{s_k, t_k\} \cap K| = 0$ then, since $d_R(s_R, t_R)$ is odd, by Lemma 2.1, there is an $s_R t_R$ -path P_R containing 2^{n-1} vertices in R . Let

$$\begin{aligned} P_i &= P'_i, \quad i = 1, 2, 3, \dots, k-1, \\ P_k &= s_k s_R + P_R + t_R t_k. \end{aligned} \quad (3.4)$$

Then $P_1, \dots, P_{k-1}, P_k$ are k disjoint fault-free ST -paths containing at least $2^n - 2f$ vertices, as required.

If $|\{s_k, t_k\} \cap K| = 2$, then there are two cases to be considered.

a) s_k and t_k are both in the same path in (3.3), say P'_1 . Let x_L and y_L be two neighbors of s_k and t_k in P'_1 but not in the subpath $P'_1(s_k, t_k)$, respectively. Since $d_L(s_k, t_k)$ is odd, both $d_L(x_L, y_L)$ and $d_R(x_R, y_R)$ are odd. By Lemma 2.1, there is an $x_R y_R$ -path P_R containing 2^{n-1} vertices in R . Let

$$\begin{aligned} P_1^* &= P'_1(s_1, x_L) + x_L x_R + P_R + y_R y_L + P'_1(y_L, t_1), \\ P_k^* &= P'_1(s_k, t_k). \end{aligned}$$

Replacing P_1 and P_k in (3.4) by P_1^* and P_k^* yields k disjoint fault-free ST -paths containing at least $2^n - 2f$ vertices.

b) s_k and t_k are in different paths in (3.3). Without loss of generality, suppose that s_k is in P'_1 and t_k is in P'_2 , two end-vertices of P'_i are s_i and t_i for $i = 1, 2$. Let x_L be the neighbor of s_k in $P'_1(s_1, s_k)$, and y_L be the neighbor of t_k in $P'_2(t_k, t_2)$. Since $d_L(s_k, t_k)$ is odd, both $d_L(x_L, y_L)$ and $d_R(x_R, y_R)$ are odd. By Lemma 2.1, there is an $x_R y_R$ -path P_R containing 2^{n-1} vertices in R . Let

$$\begin{aligned} P_1^* &= P'_1(s_1, x_L) + x_L x_R + P_R + y_R y_L + P'_2(y_L, t_2), \\ P_2^* &= P'_2(s_2, t_k), \\ P_k^* &= P'_1(s_k, t_1). \end{aligned}$$

Replacing P_1, P_2 and P_k in (3.4) by P_1^*, P_2^* and P_k^* yields k disjoint fault-free ST -paths containing at least $2^n - 2f$ vertices.

The remaining case is $|\{s_k, t_k\} \cap K| = 1$. Without loss of generality, assume $s_k \in K$ and $t_k \notin K$. We can further assume that s_k is in P'_1 with two end-vertices s_1 and t_1 . Let x_L be the neighbor of s_k in $P'_1(s_1, s_k)$, and t_R be the neighbor of t_k in R . Since $f_R = 0$ and x_R and t_R both are in X , there is a free-fault $x_R t_R$ -path P_R containing $2^{n-1} - 1$. Let

$$\begin{aligned} P_1^* &= P'_1(s_1, x_L) + x_L x_R + P_R + t_R t_k \\ P_k^* &= P'_1(s_k, t_1). \end{aligned}$$

Replacing P_1 and P_k in (3.4) by P_1^* and P_k^* yields k disjoint fault-free ST -paths containing vertices at least

$$(2^{n-1} - 2f + |\{t_k\}|) + (2^{n-1} - 1) = 2^n - 2f.$$

Summing up the above discussion, the theorem holds when $q = k$.

For sake of convenience, we assume $f_L \leq f_R$ in the following two cases.

Case 2. $1 \leq q \leq k - 1$ or $p > q = 0$ and $f_R \leq 2n - 2k - 5$.

Let W_L be the set of neighbors of vertices of S_R in L . Then $W_L \subset Y$ since $S \subset X$. For $n \geq 5$ and $k \geq 2$, we have that

$$\begin{aligned} |Y \cap V(L)| - |T_L| - |W_L| - f &= 2^{n-2} - q - (k - p) - f \\ &\geq 2^{n-2} - q - (k - p) - (2n - 2k - 3) \\ &= (2^{n-2} - 2n + k + 3) + (p - q) \\ &> p - q. \end{aligned}$$

This implies that there is a set U_L of $(p - q)$ fault-free vertices in $Y \cap V(L - T_L)$ such that its neighbor-set U_R in R is in $X \cap V(R - S_R)$ and fault-free. If $p > q$, let

$$U_L = \{u_1, u_2, \dots, u_{p-q}\} \text{ and } U_R = \{v_1, v_2, \dots, v_{p-q}\}, \quad (3.5)$$

be any two such vertex-sets, where $u_i v_i \in E(Q_n)$ for each $i = 1, 2, \dots, p - q$. If $p = q$, let $U_L = U_R = \emptyset$.

Since $f_L \leq f_R$, then $f_L \leq \frac{1}{2}f \leq \frac{1}{2}(2n - 2k - 3) \leq 2(n - 1) - 2k - 3$. By the induction hypothesis, in L there are $p(\leq k)$ disjoint fault-free paths

$$P'_1, \dots, P'_{p-q}, P_{p-q+1}, \dots, P_p$$

connecting S_L and $T_L \cup U_L$, and containing at least $2^{n-1} - 2f_L$ vertices, where P'_i connect some vertex in S_L to the vertex u_i in U_L for each $i = 1, 2, \dots, p - q$ provided that $p > q$.

If $q \geq 1$, then $f_R \leq f \leq 2n - 2k - 3 \leq 2(n - 1) - 2(k - q) - 3$ and $k - q \leq k - 1 \leq n - 3$. If $q = 0$ and $f_R \leq 2n - 2k - 5$, then $f_R \leq 2(n - 1) - 2k - 3$, $k \leq n - 3$. In any case, by the induction hypothesis, in R there are $k - q$ disjoint fault-free paths

$$P''_1, \dots, P''_{p-q}, P_{p+1}, \dots, P_k$$

connecting $S_R \cup U_R$ and T_R , and containing at least $2^{n-1} - 2f_R$ vertices, where P''_i connect the vertex v_i in U_R to some vertex in T_R for each $i = 1, 2, \dots, p - q$ provided that $p > q$. Let

$$P_i = P'_i + u_i v_i + P''_i \quad i = 1, 2, \dots, p - q.$$

Then k paths $P_1, P_2, \dots, P_{p-q}, P_{p-q+1}, \dots, P_p, P_{p+1}, \dots, P_k$ satisfy our requirements.

Case 3. $p > q = 0$ and $f_R \geq 2n - 2k - 4$.

In this case, $T_R = T$ and $f_L \leq 1$. We can write (3.5) as

$$U_L = \{u_1, u_2, \dots, u_p\} \text{ and } U_R = \{v_1, v_2, \dots, v_p\}. \quad (3.6)$$

Since $f_L \leq 1$, by induction, in L there are p disjoint fault-free $S_L U_L$ -paths

$$P'_1, P'_2, \dots, P'_p \quad (3.7)$$

containing at least $2^{n-1} - 2f_L$ vertices, without loss of generality, say P'_i connect s_i to u_i for each $i = 1, 2, \dots, p$.

Since $f_R \leq 2n - 2k - 3 = 2(n - 1) - 2(k - 1) - 3$, by the induction hypothesis, there are $k - 1$ disjoint fault-free paths

$$P''_2, \dots, P''_p, P_{p+1}, \dots, P_k \quad (3.8)$$

connecting $S_R \cup U_R - \{v_1\}$ and $T - \{t_1\}$ in R , and containing at least $2^{n-1} - 2f_R$ vertices, where P''_i connect the vertex v_i in U_R and some vertex in $T - \{t_1\}$ for each $i = 2, 3, \dots, k$. Then

$$\begin{aligned} P_1 &= P'_1, \\ P_i &= P'_i + u_i v_i + P''_i, \quad i = 2, 3, \dots, p \\ P_i &, \quad i = p + 1, \dots, k \end{aligned} \quad (3.9)$$

are k disjoint fault-free paths between S and $(T - \{t_1\}) \cup \{u_1\}$ containing at least $2^n - 2f$ vertices in Q_n . Without loss of generality, say P_i connect s_i to t_i for each $i = 2, 3, \dots, k$, P_1 connect s_1 to u_1 . Let

$$B = V(P_2 \cup \dots \cup P_k).$$

Note if $p = 1$, P'_i in (3.7) and P''_i in (3.8) are empty, and we can choose t_1 such that $s_1 t_1 \notin E(Q_n)$.

There are two subcases to be considered.

a) $t_1 \notin B$. Let t_L be the neighbor of t_1 in L .

Suppose that t_L is fault-free. If $t_L \in S$, say $t_L = s_1$. By assumption $s_1 t_1 \notin E(Q_n)$ if $p = 1$, then $p \geq 2$. We consider t_L a temporary faulty vertex. Let $S' = S - \{s_1\}$, $U' = U_L - \{u_1\}$. Then L contains at most two faulty vertices, $p - 1$ disjoint fault-free $S'U'$ -paths $P'_2, \dots, P'_p$ shown in (3.7) contain at least $2^{n-1} - 2(f_L + 1)$ vertices, $P_2, P_3, \dots, P_k$ in (3.9) are $k - 1$ disjoint paths containing at least $2^n - 2f - 2$ vertices. Together with the path $P_1 = s_1 t_1$, obtain the k paths satisfied the requirements. If $t_L \notin S$, since $f_L \leq 1$, $|F_L \cup (U_L - \{u_1\})| \leq k \leq n - 3$, thus t_L has at least a fault-free neighbor $z \in V(L) - (U_L - \{u_1\})$. Since $z \in Y$, we can choose u_1 of U_L be z in (3.6). We consider t_L a temporary faulty vertex. Then L contains at most two faulty vertices, p disjoint fault-free $S_L U_L$ -paths shown in (3.7) contain at least $2^{n-1} - 2(f_L + 1)$ vertices. Let

$$P_1^* = P_1 + u_1 t_L + t_L t_1.$$

Replacing P_1 in (3.9) by P_1^* yields k disjoint fault-free ST -paths in Q_n . Note that t_1 and t_L are two new vertices. Then k paths $P_1^*, P_2, \dots, P_k$ satisfy our requirements.

Now suppose that t_L is a faulty vertex. Then t_L is the only faulty vertex in L . Take a fault-free neighbor, say w_R of t_1 in R . Then its neighbor w_L in L is fault-free and in Y . If $w_R \notin B$, then choose $u_1 = w_L \in U_L$ in (3.6). Let

$$P_1^* = P_1 + u_1 w_R + w_R t_1.$$

Replacing P_1 in (3.9) by P_1^* yields k disjoint fault-free ST -paths in Q_n , as required. Assume that w_R is in some P_i ($2 \leq i \leq k$), say in $P_2(s_2, t_2)$. Then $w_R \neq t_2$ since w_R and t_2 are in different partite sets. Let u_R be the neighbor of w_R in $P_2(w_R, t_2)$ and u_L be the neighbor of u_R in L . Then $u_L \in X$.

If $u_L \in S$, say $u_L = s_1$. If $p = 1$, we can choose another w_R such that $u_L \neq s_1$, so let $p \geq 2$. We consider s_1 be a temporary fault vertex. Let $S' = S - \{s_1\}$, $U' = U_L - \{u_1\}$. Then L contains at most two faulty vertices, there are $p - 1$ disjoint fault-free $S'U'$ -paths $P'_2, \dots, P'_p$ containing at least $2^{n-1} - 2(f_L + 1)$ vertices, $P_2, P_3, \dots, P_k$ in (3.9) are $k - 1$ disjoint paths containing at least $2^n - 2f - 2$ vertices. Let

$$\begin{aligned} P_1 &= s_1 u_R + P_2(u_R, t_2), \\ P_2^* &= P_2(s_2, w_R) + w_R t_1. \end{aligned}$$

Thus $P_1, P_2^*, P_3, \dots, P_k$ are k disjoint fault-free ST -paths in Q_n , where t_1 and s_1 are two new vertices, as required.

If $u_L \notin S$, choose a fault-free neighbor z_L of u_L in L such that it is not any u_i in U_L for $2 \leq i \leq p$. Then $z_L \in Y$. So choose $u_1 = z_L \in U_L$ in (3.6). We consider u_L a temporary faulty vertex. Then L contains at most two faulty vertices. k disjoint fault-free SU_L -paths shown in (3.7) contain at least $2^{n-1} - 2(f_L + 1)$ vertices. Let

$$\begin{aligned} P_1^* &= P_1 + z_L u_L + u_L u_R + P_2(u_R, t_2), \\ P_2^* &= P_2(s_2, w_R) + w_R t_1. \end{aligned}$$

Replacing P_1 and P_2 in (3.9) by P_1^* and P_2^* yields k disjoint fault-free ST -paths in Q_n , where t_1 and u_L are two new vertices, as required.

b) $t_1 \in B$.

In this case, without loss of generality, we assume t_1 is in P_2 that connects s_2 to t_2 . Let u_R and v_R be two neighbors of t_1 in P_2 , and let u_L and v_L be two neighbors of u_R and v_R in L , respectively. Then u_L and v_L both are in Y . Without loss of generality, assume that v_R is in $P_2(s_2, t_1)$ and u_R is in $P_2(t_1, t_2)$.

If $f_L = 0$ or u_L is fault-free, then choose $u_1 = u_L$ in U_L in (3.6). Let

$$\begin{aligned} P_1^* &= P_2(s_2, t_1), \\ P_2^* &= P_1 + u_L u_R + P_2(u_R, t_2). \end{aligned}$$

Replacing P_1 and P_2 in (3.9) by P_1^* and P_2^* yields k disjoint fault-free ST -paths in Q_n , as required.

Assume now that $f_L = 1$ and u_L is fault vertex. Since $f \leq 2n - 2k - 3$, $f_R = 2n - 2k - 4$. If $d(u_R, t_2) > 1$, we can choose t_2 instead of t_1 , which comes down to the above case, yields k disjoint fault-free ST -paths as required. Thus, we can assume $d(u_R, t_2) = 1$.

Let t_{2L} be the neighbor of t_2 in L , then $u_L t_{2L} \in E(Q_n)$. If $t_{2L} \in S_L$, say $t_{2L} = s_1$. By assumption $s_1 t_1 \notin E(Q_n)$ if $p = 1$, then $p > 1$. Let $S' = S - \{s_1\}$, $U' = U_L - \{u_1\}$. Using Lemma 2.4 and choosing xy to be $u_L t_{2L}$, then there is $p - 1 (\leq k)$ disjoint $S'U'$ -paths

$$P'_2, \dots, P'_p \tag{3.10}$$

that contain all vertices in $L - \{u_L, t_{2L}\}$, say P'_i connect s_i to u_i for each $i = 2, 3, \dots, p$. Combining (3.8) with (3.10), we obtain $k - 1$ disjoint fault-free paths between S' and $T - \{t_1\}$

$$\begin{aligned} P_i &= P'_i + u_i v_i + P''_i, \quad i = 2, 3, \dots, p \\ P_i \quad &i = p + 1, \dots, k \end{aligned} \tag{3.11}$$

that contain at least $(2^{n-1} - 2) + (2^{n-1} - 2f_R) = 2^n - 2f$ vertices in Q_n . Let

$$P_1 = s_1 t_2, \quad \text{and} \quad P_2^* = P_2(s_2, t_1).$$

then the paths $P_1, P_2^*, P_3, \dots, P_k$ as required.

If $t_{2L} \notin S_L$, let $z \in N(t_{2L}) \cap L - \{u_L\} - U'$. Choose $u_1 = z$ in U_L in (3.6). Using Lemma 2.4 and choosing xy to be $u_L t_{2L}$, then there is $p (\leq k)$ disjoint $S_L U_L$ -paths

$$P'_1, P'_2, \dots, P'_p \tag{3.12}$$

that contain all vertices in $L - \{u_L, t_{2L}\}$, say P'_i connect s_i to u_i for each $i = 1, 2, \dots, p$. Combining (3.8) with (3.12), we obtain k disjoint fault-free paths between S and $(T - \{t_1\}) \cup \{u_1\}$

$$\begin{aligned} P_1 &= P'_1, \\ P_i &= P'_i + u_i v_i + P''_i, \quad i = 2, 3, \dots, p \\ P_i, \quad i &= p+1, \dots, k \text{ (defined in (3.8))} \end{aligned} \tag{3.13}$$

that contain at least $(2^{n-1} - 2) + (2^{n-1} - 2f_R) = 2^n - 2f$ vertices in Q_n . Let

$$P_1^* = P_1 + u_1 t_{2L} + t_{2L} t_2 \quad \text{and} \quad P_2^* = P_2(s_2, t_1).$$

Replacing P_1 and P_2 in (3.13) by P_1^* and P_2^* yields k disjoint fault-free ST -paths in Q_n . We remove one vertex u_R from paths in (3.11), and add one vertex t_{2L} to obtain new paths. Thus, these paths still contain at least $2^n - 2f$ vertices in Q_n , as required.

The theorem follows.

References

- [1] C.-H. Chang, C.-K. Lin, H.-M. Huang, L.-H. Hsu, The super laceability of the hypercubes. *Information Processing Letters*, 92 (2004), 15-21.
- [2] C.-H. Chang, C.-M. Sun, H.-M. Huang, L.-H. Hsu, On the equitable k^* -laceability of hypercubes. *Journal of Combinatorial Optimization*, 14 (2007), 349-364.
- [3] X.-B. Chen, Many-to-many disjoint paths in faulty hypercubes. *Information Sciences*, 179 (2009), 3110-3115.
- [4] T. Dvořák, Hamiltonian cycles with prescribed edges in hypercubes. *SIAM J. Discrete Math.*, 19 (2005), 135-144.
- [5] T. Dvořák, V. Koubek Long paths in hypercubes with a quadratic number of faults. *Information Sciences*, 179 (2009), 3763-3771.
- [6] J. Fink and P. Gregor. Long paths and cycles in hypercubes with faulty vertices. *Information Sciences*, 179 (2009), 3634-3644.
- [7] F. Harary, J. P. Hayes, and H.-J. Wu, A survey of the theory of hypercube graphs. *Computers and Mathematics with Applications*, 15 (4) (1988), 277-289.
- [8] P. Gregor, R. Škrekovski, Long cycles in hypercubes with distant faulty vertices. *Discret. Math. Theor. Comput. Sci.*, 11 (1) (2009), 185-198.
- [9] T.-L. Kueng, T. Liang, L.-H. Hsu, J. J. M. Tan, Long paths in hypercubes with conditional node-faults. *Information Sciences*, 179 (2009), 667-681.
- [10] F. T. Leighton, *Introduction to Parallel Algorithms and Architectures: Arrays, Trees, Hypercubes*, Morgan Kaufmann, San Mateo, CA, 1992.

- [11] M.-J. Ma, The spanning connectivity of folded hypercubes. *Information Sciences*, 180 (2010), 3373-3379.
- [12] J.-H. Park, H.-C. Kim and H.-S. Lim, Many-to-many disjoint path covers in the presence of faulty elements. *IEEE Trans. Comput.* 58 (2009), 528-540.
- [13] J.-M. Xu, *Topological Structure and Analysis of Interconnection Networks*. Kluwer Academic Publishers, Dordrecht/Boston/London, 2001.
- [14] J.-M. Xu, *Theory and Application of Graphs*. Kluwer Academic Publishers, Dordrecht/Boston/London, 2003.
- [15] J.-M. Xu, and M.-J. Ma, A survey on cycle and path embedding in some networks. *Frontiers of Mathematics in China*, 4 (2) (2009), 217-252.